

Eur J Vasc Endovasc Surg 26, 418–422 (2003)

doi: 10.1016/S1078-5884(03)00177-1, available online at <http://www.sciencedirect.com> on  SCIENCE @ DIRECT®

## Systemic Levels of Cotinine and Elastase, but not Pulmonary Function, are Associated with the Progression of Small Abdominal Aortic Aneurysms

J. S. Lindholt,<sup>1\*</sup> B. Jørgensen,<sup>2</sup> N. A. Klitgaard<sup>2</sup> and E. W. Henneberg<sup>3</sup><sup>1</sup>Department of Vascular Surgery, Rigshospitalet, University of Copenhagen, Denmark and<sup>2</sup>Departments of Clinical Biochemistry, and <sup>3</sup>Vascular Surgery, Viborg Hospital, Denmark

**Objective:** to study whether smoking and impaired pulmonary function are associated with the expansion of abdominal aortic aneurysms (AAA).

**Methods and material:** seventy-nine men with small (3–5 cm), screen-detected AAA underwent a simple 5-step smoking history, measurement of the forced first second expiratory volume (FEV1), venepuncture and annual ultrasound scan for mean follow-up period of 3.5 years.

**Results:** all but one patient had a significantly reduced FEV1 ( $p < 0.05$ , Mann–Whitney). The FEV1/expected FEV1 ratio (rFEV1) was not related to AAA expansion but was negatively correlated with P-elastase- $\alpha_1$ -antitrypsin-complexes (P-Elastase). P-Elastase was positively correlated with smoking and S-cotinine. Smoking, S-cotinine, and P-elastase were positively correlated with the mean annual AAA expansion rate but not rFEV1.

**Conclusion:** in general, patients with AAA have impaired pulmonary function. A simple five step smoking classification is as predictive of AAA-expansion as S-cotinine. Smoking may cause elastase secretion leading to pulmonary and aortic elastin degradation but the lack of association between AAA-expansion and rFEV1 suggest that other mechanisms are important.

**Key Words:** Abdominal aortic aneurysms; Surveillance; Pathogenesis; Smoking; Pulmonary function.

### Introduction

Patients with chronic obstructive pulmonary disease (COPD) are more likely to have abdominal aortic aneurysms (AAA) and AAA in such patients may be more prone to rupture.<sup>1–12</sup> The link may be related to the degradation of elastic tissue by elastase in the lungs and the aorta as a result of smoking.<sup>13–22</sup> Specifically, smoking inhibits  $\alpha_1$ -antitrypsin, the major inhibitor of elastase, and stimulates elastase secretion from the neutrophils. The U.K. small aneurysm trial showed a relationship between decreasing FEV1 and AAA rupture<sup>23</sup> and smoking is clearly associated with the development,<sup>6,9,11,24,25</sup> and possibly the expansion and rupture, of AAA.<sup>26,27</sup> The aim of the present study was to examine the relationship between smoking history, serum markers

of nicotine consumption (cotinine),<sup>28</sup> lung function, serum markers of elastolysis and AAA expansion in a series of patients with small, screen-detected AAA.

### Material and Method

The Viborg country AAA screening programme has been described previously.<sup>7</sup> Seventy-nine of all 110 small (3–5 cm), screen-detected AAA in 1994, who attended at least one annual control scanning were randomly selected to undergo a simple 5-step smoking history, measurement of the forced first second expiratory volume (FEV1),<sup>29</sup> venepuncture and annual ultrasound scan for mean follow-up of 3.5 years. The relative FEV1 (rFEV1) was defined as actual FEV1/predicted FEV1.<sup>30</sup> The mean annual expansion was calculated as: ((Present anteriorposterior (AP)-diameter—Initial AP-diameter)/days of observation)\*365.25 days. The two observers<sup>31</sup> were blind to the results from the blood-samples. S- $\alpha_1$ -antitrypsin, P-elastase- $\alpha_1$ -antitrypsin-complexes (P-Elastase) (non-commercial ELISA), and S-Cotinine (RIA, Diagnostic Products Corp., LA, U.S.A.) were determined by

The paper was presented at the American Association for Vascular Surgery 50th Scientific Meeting and The Society for Vascular Surgery 56th Annual meeting, June 9–12, 2002, at the Hynes Convention Center in Boston, MA, U.S.A.

\*Corresponding author. J. S. Lindholt, Department of Vascular Surgery RK3111, Rigshospitalet, Blegdamsvej 15, DK-2100 Copenhagen Ø.

**Table 1.** Medians and interquartiles of the various analysed parameters among those selected for the study and those who were not selected to the study.

	Selected study group <i>n</i> = 79			Non-selected group <i>n</i> = 31			<i>p</i> -value
	25th Quartiles	Median	75th Quartiles	25th Quartiles	Median	75th Quartiles	
Age (years)	67	69	71	66	68	70	0.15
AAA-size (mm)	31	33	39	30	32	38	0.40
Expansion rate (mm/year)	0.93	2.34	3.83	0.54	2.23	5.48	0.87
Systolic BP (mmHg)	140	160	170	150	160	175	0.38
Diastolic BP (mmHg)	85	90	100	90	100	100	0.43
ABI* (per cent)	81	100	114	85	100	116	0.52
rFEV1 (per cent)	52.0	72.5	91.5	53.0	77.0	86.0	0.96
Smokers (per cent)		37.5			39.2		0.87

*p*-values from a non-parametric comparison of these two subgroups are shown in the right column.

\*ABI = Ankle brachial blood pressure index.

standard methods as previously described.<sup>18,32,33</sup> Smoking habits were classified as: 0 = never or former smokers, 1 = 1–10 cigarettes daily, 2 = 11–20 cigarettes daily, 3 = above 20 cigarettes daily. All others, e.g., smokers with a mixed consumption of pipe, cigars, and cigarettes, were ranked 4 based empirically upon the median cotinine levels. The trial was approved by the local scientific ethics committee and reported to the Central Control of Registers. SPSS 10.0 was used for non parametric tests, Spearman's correlation, Mann–Whitney, and Kruskal–Wallis.<sup>34</sup>


## Results

Demographic data of the selected and unselected cases are presented in Table 1 showing no apparent differences. The median annual AAA expansion was


2.3 mm, FEV1 was 2.25 (interquartile range [IQR] 1.56–2.88), and the rFEV1 was 0.70 (IQR 0.46–0.93). All of the patients bar one had a FEV1 significantly below expected ( $p < 0.05$ , Mann–Whitney). The smoking scale correlated significant positively with S-Cotinine, P-Elastase, S- $\alpha_1$ -antitrypsine, and expansion rate but not with rFEV1 (Table 2). S-Cotinine also correlated with P-Elastase, S- $\alpha_1$ -antitrypsin, and expansion rate (Fig. 1) but not with rFEV1 (Table 3). P-Elastase also correlated negatively with rFEV1, positively with expansion rate (Fig. 2) and S- $\alpha_1$ -antitrypsin (Table 3). Finally, rFEV1 and mean annual expansion rate did not correlate ( $r = 0.13$ ,  $p = 0.26$ ) (Table 3, Fig. 3).

## Discussion

All patients bar one had impaired lung function.<sup>30</sup> The



**Fig. 1.** Scatter plot of serum level of S-Cotinine and the expansion rate of small abdominal aortic aneurysms. Spearman's correlation coefficient  $r = 0.24$  ( $p = 0.04$ ).




**Fig. 2.** Scatter plot of plasma level of P-Elastase and the expansion rate of small abdominal aortic aneurysms. Spearman's correlation coefficient  $r = 0.30$  ( $p = 0.0004$ ).

**Table 2. Comparison of smoking interview data classified into a five categorical rank scale with serological markers, pulmonary function, and the size and expansion rate of small abdominal aortic aneurysms.**

Smoking scale	N	S-Cotinine ng/ml	P-Elastase ng/ml	$\alpha_1$ -anti-trypsin mg/l	rFEV1 l/min	Expansion rate mm/year	Initial AAA-size mm
Non-smokers	31	33.1	50.7	1.26	2.31	1.13	34.7
<10 cig. Daily	12	259.9	69.8	1.50	2.33	2.00	34.9
10–20 cig. Daily	20	305.3	67.0	1.41	2.08	2.70	34.5
>20 cig. Daily	5	303.1	64.4	1.34	2.09	2.25	36.6
Mixed consumption	11	327.4	89.3	2.08	2.18	4.56	35.5
Kruskal–Wallis test	79	(0.000)	(0.001)	(0.015)	(0.818)	(0.011)	(0.861)
Correlation coefficient, $r$ ( $p$ -value)	79	0.69 (0.000)	0.46 (0.000)	0.31 (0.006)	–0.09 (0.401)	0.22 (0.048)	0.03 (0.798)

Medians with 25- and 75 percentiles. Kruskal–Wallis tests for differences between the ranks of the scale, and Spearman's correlation coefficients for correlation between the ranks.  $p$ -values in parentheses. Last row: Spearman's correlation coefficients with  $p$ -value in parentheses.



**Fig. 3.** Scatter plot of the relative pulmonary function and the expansion rate of small abdominal aortic aneurysms. Spearman's correlation coefficient  $r = 0.13$  ( $p = 0.26$ ).

negative correlation between FEV1 and P-Elastase suggest that this is, at least partly, caused by elastolysis. Similarly, the correlation between P-Elastase and expansion implicates elastolysis in aneurismal degradation. Perhaps surprisingly, therefore, there was no correlation between FEV1 and expansion. This may be because elastase plays a major role in the pulmonary disease but only a small role in AAA, where other proteases such as plasmin, metalloproteinases, and cysteine proteases are also active.<sup>35–40</sup>

As cotinine has a half-life of about 18 h (cf. 2 h for nicotine), plasma levels are constant during the smoking hours. As urinary cotinine needs to be creatinine-adjusted the measurement of plasma cotinine is preferable.<sup>41–44</sup> The simple 5-step smoking history obtained at interview was just as closely associated with AAA expansion as P-Cotinine levels, FEV1, P-Elastase, and S- $\alpha_1$ -antitrypsin levels. This reaffirms the importance of implementing effective smoking cessation strategies in this patient population.

To summarise, most men with AAA have impaired pulmonary function. Smoking and elastase-secretion are predictive for pulmonary and aortic elastin degradation but the lack of association between AAA-expansion and FEV1 suggests that additional mechanisms are at work in the matrix degradation of the aorta causing AAA.

Table 3. Correlation matrix of mean annual expansion rate, relative pulmonary function (rFEV1) and serological markers.

	Smoking	S-Cotinine	P-Elastase	rFEV1
S-Cotinine (ng/ml)	0.69 (<0.01)			
P-Elastase (ng/ml)	0.46 (<0.01)	0.48 (<0.01)		
rFEV1 (l/min)	-0.07 (0.57)	-0.14 (0.24)	-0.23 (0.05)	
$\alpha_1$ -antitrypsin (ng/ml)	0.31 (<0.01)	0.37 (<0.01)	0.28 (0.01)	-0.17 (0.15)
Expansion rate (mm/year)	0.23 (0.04)	0.24 (0.04)	0.30 (<0.01)	0.13 (0.26)

Spearman's correlation coefficients. *p* values in parentheses. rFEV1 = Observed FEV1/expected FEV1.

### Acknowledgements

Nurse A. K. Sahlholdt, Dr P. Jaeger, and Dr S. Vammen are thanked for committed assistance. The Health Department of Viborg County, The Danish Heart Foundation, The Danish National Council of Health Research, The Foundation of Health Research in Western Denmark, and The Foundation of Rosa and Asta Jensen are thanked for economical support.

### References

- 1 VAN LAARHOVEN CJHM, BORSTLAP ACW, VAN BERGE HENEGOUWEN DP *et al.* Chronic obstructive pulmonary disease and abdominal aortic aneurysms. *Eur J Vasc Surg* 1993; **7**: 386–390.
- 2 BENGTSSON H, BERGQVIST D, EKBERG O, JANZON L. A population based screening of abdominal aortic aneurysms. *Eur J Vasc Surg* 1991; **5**: 53–57.
- 3 SIMONI G, PASTORINO C, PERRONE R *et al.* Screening for abdominal aortic aneurysms and associated risk factors in a general population. *Eur J Vasc Endovasc Surg* 1995; **10**: 207–210.
- 4 JONES HJS, IBRAHIM AEK, HOSKINS C, DERODRA JK. Aortic screening in general practice. *Br J Surg* 1996; **348**: 1320.
- 5 SCOTT RA, WILSON NM, ASHTON HA, KAY DN. Influence of screening on the incidence of ruptured abdominal aortic aneurysm: 5-year results of a randomized controlled study. *Br J Surg* 1995; **82**: 1066–1070.
- 6 PLEUMEEKERS HJCM, HOES AW, VAN DER DOES E *et al.* Aneurysms of the abdominal aorta in older adults. The Rotterdam study. *Am J Epidemiol* 1995; **142**: 1291–1299.
- 7 LINDHOLT JS, HENNEBERG EW, FASTING H, JUUL S. High-risk or mass screening for abdominal aortic aneurysms. *Br J Surg* 1997; **84**: 40–42.
- 8 MECHAM RP, LANGE G. Antibodies to insoluble and solubilized elastin. *Methods in Enzymology* 1982; **82**: 744–759.
- 9 KROHN CD, KULLMANN G, KVERNEBO K, ROSEN L, KROESE A. Ultrasonographic screening for abdominal aortic aneurysm. *Eur J Surg* 1992; **158**: 527–530.
- 10 LUCAROTTI ME, SHAW E, POSKITT K, HEATHER B. The Gloucestershire Aneurysm Screening Programme: The first 2 years experience. *Eur J Vasc Surg* 1993; **7**: 397–401.
- 11 SMITH FCT, GRIMSHAW GM, PATERSON IS, SHEARMAN CP, HAMER JD. Ultrasonographic screening for abdominal aortic aneurysm in an urban community. *Br J Surg* 1993; **80**: 1406–1409.
- 12 COLLIN J, ARAUJO L, WALTON J, LINDSELL D. Oxford screening programme for abdominal aortic aneurysm in men 65–75 years. *Lancet* 1988; **318**: 613–615.
- 13 DILLON TJ, WALSH RL, SCICCHITANO R, ECKERT B, CLEARY EG, MCLENNAN G. Plasma elastin-derived levels in normal adults, children and emphysematous subjects. *Am Rev Respir Dis* 1992; **146**: 1143–1148.
- 14 SCHRIVER EE, DAVIDSON JM, SUTCLIFFE MC, SWINDELL BB, BERNARD GR. Comparison of elastin peptide concentrations in body fluids from healthy volunteers, smokers and patients with chronic obstructive pulmonary disease. *Am Rev Respir Dis* 1992; **145**: 762–766.
- 15 JANOFF A. Elastases and Emphysema. *Am Rev Respir Dis* 1985; **132**: 417–433.
- 16 FUJITA J, ABBODD RT, RICHTER A, JOHAL SHS. Evaluation of elastase and antielastase balance in patients with chronic bronchitis and pulmonary emphysema. *Am Rev Respir Dis* 1990; **142**: 57–61.
- 17 WRIGHT RR. Elastic tissue of normal and emphysematous lungs. A three-dimensional study. *Am J Pathol* 1961; **39**: 355–363.
- 18 LINDHOLT JS, HEICKENDORFF L, FASTING H, HENNEBERG EW. Serum-elastin-peptides as predictor for the expansion of small abdominal aortic aneurysms. *Eur J Vasc Endovasc Surg* 1997; **14**: 12–16.
- 19 FÜLOP JR T, WEI SM, ROBERT L, JACOB M-P. Determination of elastin peptides in normal and arteriosclerotic sera by ELISA. *Clin Physiol Biochem* 1990; **8**: 273–282.
- 20 BUSUTTIL RM. Collagenase activity of the human Aorta. *Arch Surg* 1980; **115**: 1373–1378.
- 21 COHEN JR. Altered aortic protease and antiprotease activity in patients with ruptured abdominal aortic aneurysm. *Surg Gynecol Obstet* 1987; **164**: 355–357.
- 22 DOBRIN PB. Elastolytic and collagenolytic studies of arteries. *Arch Surg* 1984; **119**: 405–409.
- 23 UK Small Aneurysm Trial Participants, Smoking, lung function and the prognosis of abdominal aortic aneurysms. *Eur J Vasc Endovasc Surg* 2000; **19**: 636–642.
- 24 FRANKS PJ, EDWARDS RJ, GREENHALGH RM, POWELL JT. Risk factors for abdominal aortic aneurysms in smokers. *Eur J Vasc Endovasc Surg* 1996; **11**: 487–492.
- 25 LOUWRENS HD, ADAMSON J, POWELL JT, GREENHALGH RM. Risk factors for atherosclerosis in men with stenosing or aneurysmal disease of the abdominal aorta. *Int Angio* 1993; **12**: 21–24.
- 26 KELLY TJ, HEATHER BP. General practice-based population screening for abdominal aortic aneurysm. *Br J Surg* 1989; **76**: 479–480.
- 27 LINDHOLT JS, HEEGAARD NH, VAMMEN S, FASTING H, HENNEBERG EW, HEICKENDORFF L. Smoking, but not lipids, lipoprotein(a) and antibodies against oxidised LDL, is correlated to the expansion of abdominal aortic aneurysms. *Eur J Vasc Endovasc Surg* 2001; **21**: 51–56.
- 28 MACSWEENEY ST, ELLIS M, WORRELL PC, GREENHALGH RM, POWELL JT. Smoking and growth rate of small abdominal aortic aneurysm. *Lancet* 1994; **344**: 651–652.
- 29 WILTSHIRE N, KENDRICK AH. Evaluation of a new electronic spirometer: the Vitalograph "Escort" spirometer. *Thorax* 1994; **49**: 175–178.
- 30 QUANJER PH, TAMMELING GJ, COTES JE, PEDERSEN OF, PESLIN R, YERNAULT JC. Lung volumes and forced ventilatory flows. Report working party: Standardization and lung function testing. *Eur Respir J* 1993; **6**: 5–40.
- 31 ANTONSEN S, WANSCHER M. An ELISA for elastase  $\alpha_1$ -protease inhibitor complexes in human plasma and serum. *Scand J Clin Lab Invest* 1993; **53**: 145–153.
- 32 PERKINS L, LIVESY JF, ESCARES EA, BELCHER JM, DUDLEY DK. High performance liquid-chromatographic method compared with a modified radioimmunoassay of Cotinine in plasma. *Clin Chemistry* 1991; **37**: 1989–1993.
- 33 LINDHOLT JS, VAMMEN S, JUUL S, FASTING H, HENNEBERG EW. The validity of ultrasonographic scanning as screening method for abdominal aortic aneurysm. *Eur J Vasc Endovasc Surg* 1999; **17**: 472–475.

- 34 ARMITAGE P, BERRY G. *Statistical Methods in Medical Research*. London: Blackwell Scientific Publications, 1994.
- 35 LINDHOLT JS, JØRGENSEN B, FASTING H, HENNEBERG EW. Plasma levels of plasmin-antiplasmin-complexes are predictive for small abdominal aortic aneurysms expanding to operation-recommendable sizes. *J Vasc Surg* 2001; **34**: 611–615.
- 36 NEWMAN KM, JEAN-CLAUDE J, LI H *et al.* Cellular localisation of matrix metalloproteinases in the abdominal aortic aneurismal wall. *J Vasc Surg* 1994; **20**: 814–820.
- 37 SAKALIHASAN N, DELVENNE P, NUSGENS BV, LIMET R, LAPIÈRE CM. Activated forms of MMP2 and MMP9 in abdominal aortic aneurysms. *J Vasc Surg* 1996; **24**: 127–133.
- 38 PATEL MI, MELROSE J, GHOSH P, APPLEBERG M. Increased synthesis of matrix metalloproteinases by aortic smooth muscle cells is implicated in the etiopathogenesis of abdominal aortic aneurysms. *J Vasc Surg* 1996; **24**: 82–92.
- 39 OKADA Y, KATSUDA S, NAKANISHI I. An elastolytic enzyme detected in the culture medium of human arterial smooth cells. *Cell Bio Int* 1993; **17**: 863–969.
- 40 WILLS A, THOMPSON MM, CROWTHER M *et al.* Elastase-induced matrix degeneration in arterial organ cultures: An vitro model of aneurysmal disease. *J Vasc Surg* 1996; **24**: 667–679.
- 41 KYEREMATEN GA, DAMIANO MD, DVORCHIK BH, VESELL ES. Smoking-induced changes in nicotine disposition: application of a new HPLC assay for nicotine and its metabolites. *Clin Pharmacol Ther* 1982; **32**: 769–778.
- 42 BENOWITZ NL, KUYT F, JACOB P, JONES RT, OSMAN A-L. Cotinine disposition and effects. *Clin Pharmacol Ther* 1983; **34**: 604–611.
- 43 CHEN GF, TANG YM, GREEN B *et al.* Low frequency of CYP2A6 gene polymorphisms as revealed by a one-step polymerase chain reaction method. *Pharmacogenetics* 1999; **9**: 327–332.
- 44 BOURIAN M, GULLSTEN H, LEGRUM W. Genetic polymorphism of CYP2A6 in the German population. *Toxicology* 2000; **144**: 129–137.

Accepted 24 March 2003